Department of Veterans Affairs

(Authority: 38 U.S.C. 1155)

[29 FR 6718, May 22, 1964, as amended at 34 FR 5063, Mar. 11, 1969; 40 FR 42540, Sept. 15, 1975; 41 FR 11301, Mar. 18, 1976; 66 FR 29488, May 31, 2001]

THE GENITOURINARY SYSTEM

§4.115 Nephritis.

Albuminuria alone is not nephritis, nor will the presence of transient albumin and casts following acute febrile illness be taken as nephritis. The glomerular type of nephritis is usually preceded by or associated with severe infectious disease; the onset is sudden, and the course marked by red blood cells, salt retention, and edema; it may clear up entirely or progress to a chronic condition. The nephrosclerotic type, originating in hypertension or arteriosclerosis, develops slowly, with minimum laboratory findings, and is associated with natural progress. Separate ratings are not to be assigned for disability from disease of the heart and any form of nephritis, on account of the close interrelationships of cardiovascular disabilities. If, however, absence of a kidney is the sole renal disability, even if removal was required because of nephritis, the absent kidney and any hypertension or heart disease will be separately rated. Also, in the event that chronic renal disease has progressed to the point where regular dialysis is required, any coexisting hypertension or heart disease will be separately rated.

 $[41\ FR\ 34258,\ Aug.\ 13,\ 1976,\ as\ amended\ at\ 59\ FR\ 2527,\ Jan.\ 18,\ 1994]$

§4.115a Ratings of the genitourinary system—dysfunctions.

Diseases of the genitourinary system generally result in disabilities related to renal or voiding dysfunctions, infections, or a combination of these. The following section provides descriptions of various levels of disability in each of these symptom areas. Where diagnostic codes refer the decisionmaker to these specific areas dysfunction, only the predominant area of dysfunction shall be considered for rating purposes. Since the areas of dysfunction described below do not cover all symptoms resulting from genitourinary diseases, specific diagnoses may include a

description of symptoms assigned to that diagnosis.

Re

iat diagnosis.	
	Rat- ing
nal dysfunction:	
Requiring regular dialysis, or precluding more than sedentary activity from one of the fol-	
than sedentary activity from one of the fol-	
lowing: persistent edema and albuminuria; or,	
BUN more than 80mg%; or, creatinine more	
than 8mg%; or, markedly decreased function	
of kidney or other organ systems, estpecially	400
cardiovascular	100
Persistent edema and albuminuria with BUN 40 to 80mg%; or, creatinine 4 to 8mg%; or, gen-	
eralized poor health characterized by lethargy,	
weakness anorexia weight loss or limitation	
weakness, anorexia, weight loss, or limitation of exertion	80
Constant albuminuria with some edema; or, defi-	
nite decrease in kidney function; or, hyper-	
tension at least 40 percent disabling under di-	
agnostic code 7101	60
Albumin constant or recurring with hyaline and	
granular casts or red blood cells; or, transient	
or slight edema or hypertension at least 10	20
percent disabling under diagnostic code 7101 Albumin and casts with history of acute nephritis;	30
or, hypertension non-compensable under diag-	
nostic code 7101	0
iding dysfunction:	
Rate particular condition as urine leakage, fre-	
quency, or obstructed voiding	
Continual Urine Leakage, Post Surgical Urinary	
Diversion, Urinary Incontinence, or Stress In-	
continence:	
Requiring the use of an appliance or the wearing	
of absorbent materials which must be changed	60
more than 4 times per day Requiring the wearing of absorbent materials	60
which must be changed 2 to 4 times per day	40
Requiring the wearing of absorbent materials	
which must be changed less than 2 times per	
day	20
nary frequency:	
Daytime voiding interval less than one hour, or;	
awakening to void five or more times per night	40
Daytime voiding interval between one and two	
hours, or; awakening to void three to four	20
times per night Daytime voiding interval between two and three	20
hours, or; awakening to void two times per	
night	10
structed voiding:	
Urinary retention requiring intermittent or contin-	
uous catheterization	30
Marked obstructive symptomatology (hesitancy,	
slow or weak stream, decreased force of	
stream) with any one or combination of the fol-	
lowing: 1. Post void residuals greater than 150 cc.	
Uroflowmetry; markedly diminished peak	
flow rate (less than 10 cc/sec).	
Recurrent urinary tract infections sec-	
ondary to obstruction.	
Stricture disease requiring periodic dilata-	
tion every 2 to 3 months	10
Obstructive symptomatology with or without stric-	
ture disease requiring dilatation 1 to 2 times	
per year	0
ninary tract infection:	
Poor renal function: Rate as renal dysfunction.	
Recurrent symptomatic infection requiring drain-	
age/frequent hospitalization (greater than two	

times/year), and/or requiring continuous inten-

30

sive management ..

Urr

Oh

Rat-

§4.115b

	Rat- ing
Long-term drug therapy, 1–2 hospitalizations per year and/or requiring intermittent intensive management	10

 $[59~{\rm FR}~2527,~{\rm Jan.}~18,~1994;~59~{\rm FR}~10676,~{\rm Mar.}~7,~1994]$

§4.115b Ratings of the genitourinary

system—diagnoses.		
		Rat- ing
	Note: When evaluating any claim involving loss or loss of use of one or more creative organs, refer to § 3.350 of this chapter to determine whether the veteran may be entitled to special monthly compensation. Footnotes in the schedule indicate conditions which potentially establish entitlement to special monthly compensation; however, there are other conditions in this section which under certain circumstances also establish entitlement to special monthly compensation.	
7500	Kidney, removal of one: Minimum evaluation Or rate as renal dysfunction if there is nephritis, infection, or pathology of the other.	30
7501	Kidney, abscess of: Rate as urinary tract infection	
7502	Nephritis, chronic: Rate as renal dysfunction.	
7504	Pyelonephritis, chronic: Rate as renal dysfunction or urinary tract infection, whichever is pre- dominant.	
7505	Kidney, tuberculosis of: Rate in accordance with §§ 4.88b or 4.89, whichever is appropriate.	
7507	Nephrosclerosis, arteriolar: Rate according to predominant symptoms as renal dysfunction, hypertension or heart disease. If rated under the cardiovascular schedule, however, the percentage rating which would otherwise be assigned	

will be elevated to the next higher evaluation.

Rate as hydronephrosis, except for recurrent stone formation requiring one or more of the following:

3. invasive or non-invasive procedures more than two times/year

7508 Nephrolithiasis:

1. diet therapy 2. drug therapy

		ing
7509	Hydronephrosis: Severe; Rate as renal dysfunction.	
F	requent attacks of colic with infection (pyonephrosis), kidney function im-	
F	pairedrequent attacks of colic, requiring cath-	30
О	eter drainage Only an occasional attack of colic, not	20
7510	infected and not requiring catheter drainage	10
7510	Rate as hydronephrosis, except for recurrent stone formation requiring one or more of the following: 1. diet therapy	
7511	drug therapy invasive or non-invasive procedures more than two times/year Ureter, stricture of:	30
7011	Rate as hydronephrosis, except for recurrent stone formation requiring one or more of the following: 1. diet therapy	
	drug therapy invasive or non-invasive procedures more than two times/year	30
	Cystitis, chronic, includes interstitial I all etiologies, infectious and non-in-ious:	
7515 inte	Rate as voiding dysfunction. Bladder, calculus in, with symptoms rfering with function:	
7516	Rate as voiding dysfunction Bladder, fistula of: Rate as voiding dysfunction or urinary tract infection, whichever is pre- dominant.	
7517	Postoperative, suprapubic cystotomy Bladder, injury of: Rate as voiding dysfunction.	100
7518	Urethra, stricture of: Rate as voiding dysfunction.	
7519	Urethra, fistual of: Rate as voiding dysfunction.	
7520	Multiple urethroperineal fistulae Penis, removal of half or more Or rate as voiding dysfunction.	100 30
7521	Penis removal of glans Or rate as voiding dysfunction.	20
7522 pow	Penis, deformity, with loss of erectile ver—20 1.	
7523	Testis, atrophy complete:. Both—20 ¹ One—0 ¹	
7524	Testis, removal:.	